

Aminolean®

Sterile Injection

Amino acids, B-complex vitamins and Electrolytes

DIN 00506990



INDICATIONS

As an aid in the supportive treatment of debilitated animals and for supportive treatment of severe diarrhea in cattle, swine and horses. Aminolean aids in counteracting the protein, electrolyte and vitamin loss associated with these conditions.

DESCRIPTION

Sterile injection containing amino acids, B-complex vitamins and electrolytes for cattle, horses and swine.

PACKAGING

ITEM NO.	UNIT PACKAGE	CASE SIZE
1AM1001	500 mL	12
1AM1002	950 mL	12

See reverse side for Administration and Dosage.

Aminolean[®]

Sterile Injection

Amino acids, B-complex vitamins and Electrolytes

DIN 00506990

Veterinary Use Only

ACTIVE INGREDIENTS:

Each 100 mL contains:

Dextrose 5 g, calcium chloride dihydrate 15 mg, potassium chloride 20 mg, magnesium sulfate trihydrate 20 mg, sodium acetate trihydrate 250 mg, thiamine HCl 10 mg, nicotinamide 150 mg, pyridoxine HCl 10 mg, vitamin B12 5 mcg, riboflavin 5-phosphate sodium 4 mg, d-panthenol 5 mg, casein hydrolysate*.

Each litre of solution will provide approximately:

Dextrose 252 mmol, sodium 18 mEq, calcium 2 mEq, potassium 3 mEq, magnesium 2 mEq, acetate 18 mEq, chloride 5 mEq, sulfate 2 mEq.

***Approximate amino acid profile after hydrolysis per 100 mL:**

Alanine 0.8 mg, arginine 1 mg, aspartic acid 2 mg, glutamic acid 6.7 mg, glycine 0.6 mg, histidine 0.9 mg, isoleucine 1.6 mg, leucine 2.5 mg, lysine 2.5 mg, methionine 0.3 mg, phenylalanine 1 mg, proline 3.3 mg, serine 1.6 mg, threonine 1.4 mg, tryptophane 0.3 mg, tyrosine 0.2 mg, valine 2 mg.

INDICATIONS:

As an aid in the supportive treatment of debilitated animals and for supportive treatment of severe diarrhea in cattle, swine and horses. Aminolean aids in counteracting the protein, electrolyte and vitamin loss associated with these conditions.

DOSAGE AND ADMINISTRATION:

Warm to body temperature prior to administration. Administer slowly by intravenous or intraperitoneal injection in cattle and swine. Administer by intravenous injection **ONLY** in horses.

Mature Cattle, Horses & Swine: 2 mL/kg of body weight.

Calves, Foals & Piglets: 5 mL/kg of body weight.

Repeat if necessary as directed by a veterinarian.

CAUTIONS:

Nausea and distress may sometimes occur if this solution is administered too rapidly. Should these symptoms occur, discontinue injection until animal returns to normal. Injection may then be continued at a slower rate. Supportive treatment only. Ensure proper medical treatment for infective and pathological conditions. Do not retain partially used bottle. Contains no preservatives. Do not use unless solution is clear.

STORAGE: Store below 25°C. KEEP FROM FREEZING.

WARNING: Keep out of reach of children.

1AM1001/1AM1002
8AM1011/8AM1015