

Exodus® Double Dose (pyrantel pamoate) Paste

Equine Anthelmintic

DIN 02246767



INDICATIONS

For the treatment of parasitic infections caused by the adult stage of large strongyles (*Strongylus vulgaris*, *S. equinus*, *S. edentatus*, *Triodontophorus* spp.), small strongyles (*Cyathostomum* spp., *Cylicostephanus* spp.), pinworms (*Oxyuris equi*), ascarids (*Parascaris equorum*) and the common tapeworm (*Anoplocephala perfoliata*) in horses, ponies and foals over eight weeks of age.

See reverse side for Administration and Dosage.



BENEFITS

- **Twice the doses** of other single dose brands.
- Contains proven **pyrantel pamoate** for safe, positive worm removal
- Over 90% effective against most major equine worms
- **Apple flavoured** for palatability
- Available in easy-to-use Multi Dose, 47.2 g syringes
- Pre-measured plastic syringes provide convenient, accurate dosing
- **Health Canada Approved**

PACKAGING

ITEM NO.	UNIT PACKAGE	CASE SIZE
1EX0003	12 x 23.6 g	3
1EX0004	6 x 47.2 g	3
1EX0011	50 x 23.6 g	1

Exodus® Double Dose (pyrantel pamoate) Paste

Equine Anthelmintic

DIN 02246767

DESCRIPTION: Exodus* for deworming horses is a pale yellow paste containing pyrantel pamoate in an apple-flavoured base. Pyrantel pamoate is a compound belonging to the tetrahydropyrimidine family. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid with 34.6% base activity.

Each 47.2 g multi-dose syringe contains the equivalent of 7.2 g pyrantel base sufficient to treat two 544 kg horses or several smaller horses or ponies.

INDICATIONS: For the treatment of parasitic infections caused by the adult stage of large strongyles (*Strongylus vulgaris*, *S. equinus*, *S. edentatus*, *Triodontophorus* spp.), small strongyles (*Cyathostomum* spp., *Cylicostephanus* spp.), pinworms (*Oxyuris equi*), ascarids (*Parascaris equorum*) and the common tapeworm (*Anoplocephala perfoliata*) in horses, ponies and foals over eight weeks of age.

ADMINISTRATION: Exodus* is administered orally. Fasting prior to or following treatment is not required.

The following method of administration is recommended:

1. According to the weight of the horse set the dose ring at the correct position on the weight-scale marked on the plunger.
2. Remove the cap from the syringe.
3. The paste is best distributed on the upper surface of the tongue. Introduce the nozzle end of the syringe at the corner of the mouth (through the interdental space). Direct the syringe backwards and depress the plunger to deposit the paste onto the tongue. Raise the horse's head by a hand under the chin to assist in the swallowing process.
4. If only part of the paste has been used (see Cautions), replace the cap on the syringe nozzle and store in a cool dry place.

DOSAGE: Administer at the rate of 6.6 mg pyrantel base per kg of body weight for infections of roundworm (nematodes) and 13.2 mg pyrantel base per kg body weight for infections of tapeworms (cestodes). This dosage can be achieved by administering the product according to the following schedule:

Body Weight Range	Dose (syringes)		Position of Dose Ring on Plunger	
	Round-worms	Tapeworms	Round-worms	Tapeworms
135 to 270 kg	1/4	1/2	270 kg	540 kg
271 to 540 kg	1/2	1	540 kg	1 full syringe

Horses maintained on premises where re-infection is likely to occur, may require re-treatment.

For most effective results, re-treat these horses every 4 to 6 weeks.

CAUTIONS: The same syringe should only be used for more than one horse if the animals are healthy and are running together or are on the same premises and in direct contact with each other. It is recommended that severely debilitated animals not be treated with this drug.

WARNINGS: Do not use in horses that are to be slaughtered for use in food. Keep out of reach of children.

STORAGE: Store at a temperature between 15°C and 30°C. Protect from direct sunlight and from freezing (0°C).

Exodus is a Registered Trademark of Bimeda, Inc.