

# LincoMed<sup>®</sup> 100

## Antibiotic Injection for Swine

*(Lincomycin Injection, USP)*

DIN 0242142

**VETERINARY USE ONLY**  
**STERILE**

### DESCRIPTION

LincoMed<sup>®</sup> contains lincomycin hydrochloride, an antibiotic produced by *Streptomyces lincolnensis* var. *lincolnensis*, which is chemically distinct from all other clinically available antibiotics and is isolated as a white crystalline solid.

### INDICATIONS

LincoMed<sup>®</sup> is indicated for the treatment of infectious forms of arthritis caused by organisms sensitive to its activity. This includes most of the organisms responsible for the various infectious arthritides in swine, such as *staphylococci*, *streptococci*, *Erysipelothrix* and *Mycoplasma* spp.

It is also indicated for the treatment of swine dysentery.



### PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1LIN021	100 mL	12

### WARNING

Treated animals must not be slaughtered for use in food for at least 48 hours after the latest treatment with this drug. Keep out of reach of children.

**Bimeda - MTC Animal Health, Inc.**

420 Beavertdale Road, Cambridge, Ontario N3C 2W4

Tel. (519) 654-8000 • Fax. (519) 654-8001

sales@bimedamtc.com

See reverse side for Administration and Dosage.



[www.BimedaMTC.com](http://www.BimedaMTC.com)

## LincoMed® 100 (Lincomycin Injection, USP)

VETERINARY USE ONLY  
STERILE

DIN 02424142

**FOR INTRAMUSCULAR USE IN SWINE.  
FOR USE IN ANIMALS ONLY. NOT FOR HUMAN USE.  
KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN.**

### ACTIVE INGREDIENT:

Lincomycin (as lincomycin hydrochloride), 100 mg/mL

**PRESERVATIVE:** Benzyl alcohol, 9.45 mg/mL

### INDICATIONS:

For treatment of infectious arthritis in swine caused by susceptible strains of *Staphylococcus* spp., *Streptococcus* spp., *Erysipelothrix* spp., and *Mycoplasma* spp., and for the treatment of swine dysentery.

### DOSAGE AND ADMINISTRATION:

Administer intramuscularly, 1 mL per 10 kg body weight once daily at first signs of infection and continue treatment for 3 to 7 days.

### CAUTIONS:

Use only in swine. If no improvement is noted within 48 hours, consult a veterinarian. Loose stools following treatment have been rarely reported but the diarrhea is usually mild and transient. It is not recommended that LincoMed® 100 be used to treat or control outbreaks of swine dysentery. This is more readily achieved through the drinking water or feed since all animals in a barn must be treated. The use of LincoMed® 100 in extremely sick or moribund pigs suffering from swine dysentery is recommended until appetite and/or water consumption improvements allow adequate intake of oral medication.

### WARNINGS:

Treated animals must not be slaughtered for use in food for at least 48 hours after the latest treatment with this drug. Keep out of reach of children.

### STORAGE:

Store between 15°C and 25°C. Protect from freezing. Use within 28 days following withdrawal of first dose.

**PRESENTATION:** 100 mL vials

Manufactured by:  
Bimeda-MTC Animal Health Inc.  
Cambridge, ON N3C 2W4  
[www.bimeda.com](http://www.bimeda.com)