

# Sulfa 25% Solution

## Sodium Sulfamethazine 25% Solution

DIN 00601837



### INDICATIONS

**Cattle & Sheep:** For the treatment of foot rot, shipping fever, bacillary enteritis, metritis, bacterial scours, mastitis, bacterial respiratory diseases and for coccidiosis in sheep.

**Calves:** For the treatment of secondary infections in white scours, septicemia and prevention of coccidiosis. As an aid in the treatment of bacterial respiratory diseases.

**Swine:** For the treatment of bacterial enteritis, mastitis, metritis, and bacterial respiratory diseases.

**Horses:** For the treatment of strangles, navel ill, joint ill, bacterial enteritis and many secondary bacterial infections associated with respiratory virus infections.

**Poultry:** For the treatment of caecal coccidiosis in chickens.

See reverse side for Administration and Dosage.

### DESCRIPTION

Oral sodium sulfamethazine 25% to be used as an aid in the treatment of conditions caused by bacteria sensitive to sulfamethazine.

### PACKAGING

ITEM NO.	UNIT PACKAGE	CASE SIZE
1SUL002	4 L	12

# Sulfa 25% Solution

## Sodium Sulfamethazine 25% Solution

DIN 00601837

### *Veterinary Use Only*

#### **DESCRIPTION:**

A non-sterile solution containing 25 grams of sodium sulfamethazine in each 100 mL.

#### **INDICATIONS:**

**Cattle & Sheep:** For the treatment of foot rot, shipping fever, bacillary enteritis, metritis, bacterial scours, mastitis, bacterial respiratory diseases and for coccidiosis in sheep.

**Calves:** For the treatment of secondary infections in white scours, septicemia and prevention of coccidiosis. As an aid in the treatment of bacterial respiratory diseases.

**Swine:** For the treatment of bacterial enteritis, mastitis, metritis, and bacterial respiratory diseases.

**Horses:** For the treatment of strangles, navel ill, joint ill, bacterial enteritis and many secondary bacterial infections associated with respiratory virus infections.

**Poultry:** For the treatment of caecal coccidiosis in chickens.

#### **DOSAGE:**

**Livestock:** First day: 45 mL of 25% sodium sulfamethazine for each 50 kg of body weight. Second and following days: one half of the above dosage. Do not treat for more than 5 days.

**Beef Cattle:** Add 80 mL to each 9 litres of drinking water for the first day. Second and following days : one half the above dosage. Do not treat for more than 5 days.

**Poultry:** Mix 35 mL of product per 9 litres of drinking water for 2 days - then reduce dosage to 17.5 mL of product per 9 litres of drinking water for 5 additional days. Only allow access to medicated water.

#### **CONTRAINDICATION:**

This product must not be used in laying birds.

#### **WARNING:**

Milk taken from treated animals during treatment and within 96 hours after the latest treatment must not be used as food. Treated animals must not be slaughtered for use in food for at least 12 days after the latest treatment with this drug. This product must not be added to swine feeds.

#### **STORAGE:**

Store at room temperature.

1SUL002/8SUL204